




Time-Dependent Analysis of $B \rightarrow J/\psi K_L^0$ Decays with the BaBar Detector

Bryan Dahmes

University of California, Santa Barbara

For the BaBar Collaboration




APS Meeting
Albuquerque, 21 April 2002

Why $B \rightarrow J/\psi K_L^0$?

- The Standard Model with 3 quark generations **predicts CP Violation**

- **Complex phase** in the quark mixing matrix provides the mechanism for CP Violation
- **Unitarity constraint** can be represented graphically in the complex plane




- $B \rightarrow J/\psi K^0$ (with K^0 mixing) provides a theoretically clean means of measuring CP violation in the B system:

$$A_{CP}(t) = -\eta_{CP} \sin 2\beta \sin(\Delta m \Delta t)$$

- Advantages of $B \rightarrow J/\psi K_L^0$:
 - Most experimentally accessible mode with $\eta_{CP} = 1$
 - $B \rightarrow J/\psi K_L^0$ is the 2nd most sensitive measurement of $\sin 2\beta$ after $B \rightarrow J/\psi K_S^0$
 - Same branching fraction as $B \rightarrow J/\psi K_S^0$
 - Less efficient signal reconstruction


K_L^0 reconstruction at BaBar



- Mean life $\tau \sim 52$ nsec
- Look for a **hadronic interaction** in the detector
 - **ElectroMagnetic Calorimeter (EMC)**
 - Significant energy deposit
 - Reject π^0 photons
 - **Instrumented Flux Return (IFR)**
 - At least two layers hit
 - Reject contributions from beam backgrounds
- $E(K_L)$ not measured due to vague K_L signature
- Calculate the K_L momentum from its **direction** and B-mass constraint

$$m_B^2 = \left(E_{J/\psi} + \sqrt{m_{K_L}^2 + p_{K_L}^2} \right)^2 - \left(\vec{p}_{J/\psi} + p_{K_L} \vec{d}_{K_L} \right)^2$$


Backgrounds in $B \rightarrow J/\psi K_L$



- Background from
 - Photons (in the calorimeter)
 - K_L from $B \rightarrow J/\psi X$ modes
- Reject background events with a missing momentum requirement
 - **Calculate** missing transverse momentum in the event
 - **Project** along K_L direction
 - **Subtract** K_L transverse momentum calculated assuming $B \rightarrow J/\psi K_L$ kinematics
- **Background events** have lower missing momentum than **signal events**
- Photon background mostly eliminated
- Cut has been optimized for minimal statistical error on $\sin 2\beta$

Event Selection Summary

- Maximum likelihood fit to data
 - **Signal** and **Inclusive J/ψ** shapes taken from Monte Carlo
 - **Non- ψ background** shape and normalization taken from J/ψ mass sidebands
- Data is broken up into blocks of different purity to increase signal sensitivity
- Reject some specific $B \rightarrow J/\psi X$ modes
- 56 fb^{-1} in $\sin 2\beta$ sample
 - Reconstruct nearly 12 $J/\psi K_L$ events/ fb^{-1} (6 in the EMC, 6 in the IFR)
- Most background from real K_L from $B \rightarrow J/\psi X$ decays



Fitting for $\sin 2\beta$ in $B \rightarrow J/\psi K_L$


- The $B \rightarrow J/\psi K_L$ decay mode has **non-negligible backgrounds**
- The $\sin 2\beta$ analysis for $B \rightarrow J/\psi K_L$ must properly model the Δt and CP properties of the background
 - Itemize the background components
 - **Inclusive J/ψ events** : separate the largest contributors
 - **Fake J/ψ events**
 - Include a term for each in the likelihood fit
 - Tagging dilutions and Δt distribution taken from Monte Carlo and data sidebands
 - Monte Carlo used to validate assumptions

	EMC K_L	IFR K_L	h_{cp}
Signal	$50 \pm 4 \%$	$65 \pm 4 \%$	+ 1
J/y K^{*+} ($K_L p^+$)	11 %	12 %	0
J/y K^{*0} ($K_L p^0$)	8 %	7 %	-0.68 ± 0.07
J/y K_S ($p^0 p^0$)	3 %	1 %	- 1
Other J/y	17 %	9 %	0 ± 0.25
Non-J/y	11 %	6 %	0 ± 0.25

Overlapping γ
From π^0 fake K_L .

Jumble of modes
each < 2 %.

CP Asymmetry in $B \rightarrow J/\psi K_L^0$



- Time-Dependent CP Asymmetry given by

$$A_{CP} = \frac{N(B^0 \rightarrow f_{cp}) - N(\bar{B}^0 \rightarrow f_{cp})}{N(B^0 \rightarrow f_{cp}) + N(\bar{B}^0 \rightarrow f_{cp})}$$

$$A_{CP}(t) = -\eta_{cp} \cdot \sin 2\beta \cdot \sin(\Delta m \Delta t)$$

- Preliminary Result

$$\sin 2\beta = 0.72 \pm 0.19(\text{stat}) \pm 0.06(\text{syst})$$

- Our measurement is still dominated by statistics




Systematic Errors in $B \rightarrow J/\psi K_L$

- Systematic error for the $\sin 2\beta$ fit calculated for
 - $B \rightarrow J/\psi K_L$ events only
 - K_L + all other modes (Global fit)
- Systematic errors are also limited by statistics
- Expect systematic errors to drop as more data is accumulated


Source	K_L Only	Global
Sample Composition	0.037	0.008
$J/\psi X$ fractions	0.033	0.007
MC corrections	0.026	0.006
Assumed CP of background	0.014	0.003
Δm_d and τ_B (PDG 2000)	0.014	-
Non- ψ BG Δt	0.002	0.0004
Total	0.059	0.013

Combined fit to $\sin 2\beta$




Combined Result (preliminary):
 $\sin 2\beta = 0.75 \pm 0.09(\text{stat}) \pm 0.04(\text{syst})$


$\sin 2\beta$ by decay mode




sin2β at BaBar



One solution for b is consistent with other measurements which constrain the value of r and h



Where do we go from here?



- These results refine the initial observation of CP Violation in the B system

- $\sin 2\beta$ measured at BaBar (hep-ex/0203007)

PRELIMINARY

$$\begin{aligned} \sin 2\beta (\eta_{CP} = -1) &= 0.76 \pm 0.10 \pm 0.04 \\ \sin 2\beta (K_L) &= 0.72 \pm 0.19 \pm 0.06 \\ \sin 2\beta (\text{all}) &= 0.75 \pm 0.09 \pm 0.04 \end{aligned}$$

- K_L is the 2nd most significant mode at BaBar
- Expect 100 fb^{-1} for summer 2002
 - $\eta_{CP} = -1$ modes : $0.10 \rightarrow 0.07$
 - K_L : $0.19 \rightarrow 0.14$